

Trade Name:

Natural cardio tonic **KYUSHIN****Features**

Palpitation and **panting** are caused by various causes such as a decrease in functions of the circulatory and respiratory system, excessive tension and stress, menopause, disorder of autonomic nerve caused by cold and heat, overwork and lack of sleep, excessive smoking and alcohol, obesity, hard exercise, and so on. **Palpitation** and **panting** are also caused easily by an age related decline in bodily functions. Moreover, if you leave those symptoms behind, that will be a trigger to cause various illnesses that gradually causes a decrease in your quality of life.

KYUSHIN shows the excellent efficacy on the palpitation and panting to improve the circulation by each merit in nine kinds of natural drugs originating in animal and plants. Furthermore, restorative by **KYUSHIN** is effective when person feels the decline of energy by listlessness of the body, feels the decline of consciousness by the heat etc., and has vertigo and dizziness.

Ingredients Daily Dose

A daily dose of Kyushin (6 pills) contains 5mg of Toad Cake, 4mg of Oriental Bezoar, 5mg of Velvet Horn, 25mg of Ginseng, 6mg of Saiga Antelope Horn, 7.5mg of Pearl, 3mg of Agalloch, 2.7mg of Borneo Camphor, 8mg of Animal Bile (Hog Bile).

Manufacturer	Kyushin Pharmaceutical Co., Ltd.				
Dosage Form	Dark brown pills				
Item	30 pills	60 pills	120 pills	310 pills	630 pills
Suggested retail price (tax-excluded)	¥2,200	¥4,100	¥7,600	¥17,000	¥31,000
Japan Article Number	4987061015739	4987061015746	4987061015753	4987061015760	4987061015777
Expiry Date	5 years				

Indications

Palpitation, panting, restorative.

Directions

Adults (15 years and over): take 2 pills in each 3 times per day (morning, evening and before going to bed) with water or lukewarm water.

Take this medicine with water or lukewarm water without leaving in a mouth or chewing. (If you leave this medicine in a mouth or chew it, numbness of tongue or mouth is left for a while because of the property of an ingredient.)

**PRECAUTIONS FOR USE:**

DON' T DO

[If the following is not kept, a present symptom takes a turn for the worse and it becomes easy for the side effect to happen.]



Don' t take the following medicines while taking this medicine:

Any other cardio tonics

CONSULT:

1. The following persons should consult with a physician, pharmacist or tourokuhambaisya (registered retail person) before taking this medicine;

1. Persons who are under treatment by a physician.
2. Persons who are pregnant or may be pregnant.

2. In case that the following symptoms appear after taking this medicine, discontinue taking this medicine at once for the possibility of the side effect, and consult a physician, pharmacist or tourokuhambaisya (registered retail person) with this manual.

Areas concerned : Symptoms

Skin : Rash, redness, itching

Digestive organs: Nausea, vomiting

3. In case that the symptoms have not improved even if taking this medicine for 5~6 days, discontinue taking this medicine, and consult a physician, pharmacist or tourokuhambaisya (registered retail person) with this manual.