Name	Product Discription	Efficacy	Dosage and Administration		Ingredients		Warning	other information	
Direct Lozenge Mango R	only lozenge which contains the crude medicine. When you have a sore or irritable throat from yelling or other causes, just put	e throat pain, and other uncomfortable throat symptoms.	Suggested Usage: Take the following doses without water. Allow it to dissolve in your mouth without chewing. Have at least 2 hours between each dose.					pharmacist, or registered sales clerks. (1) Undergoing treatment for a medical	away from a direct sun light. (2) Keep out of reach of children. (3) Store medications only in their
			Age	Dosage	Daily usage	Ingredients	Contenst		original containers in order to avoid accidental ingestion.
			Adult(older than 15 years)	1tablet		Powdered Platycodon	70.0mg		(4) Do not take expired medications.
			5 ∼ under 15 years	1/2 tablet	3∼6 times	Apricot Kernel	5.0mg		
			under 5 years	Do no	ot use	Powdered Segana	3.0mg	2. If you have the following reactions to the	
			<dosage administration.="" and=""> <ol> <li>Follow the usage and dosage instruction.</li> </ol></dosage>			Powdered Grycyrrhiza Additives: Anhydrous		product, discontinue use immediately. Take this product instruction to consult with your physician, pharmacist, or registered	
			Children should have p medicine.     S.Do not crush, or swallov		on when taking	citric acid, D-sorbitol, S Mg., gelatin, glycerol, ril menthol, Flavoring	Stearic acid boflavin, L- blet are from	sales clerks. Symptoms Skin: Rash, itchiness, skin reddening Digestive system: nausea, vomiting, loss of appetite, Nervous system: Dizziness  3. If your condition is severe and persists for more than 5to 6days, please stop taking the product. Take this product instruction to consult with your physician, pharmacist, or registered sales clerks.	