

*Please store this instruction leaflet in a safe place for later reference.*

*Be sure to read the instructions carefully before use.*

## HIYA KIOGAN GINTSUBU (SILVER)

### 樋屋奇応丸 銀粒

販売名：小児薬樋屋奇応丸

“HIYA KIOGAN GINTSUBU”, containing 4 kinds of highly-selected crude drug, mildly improves various symptoms of children, such as night crying, loss of appetite, *etc.*

### CAUTIONS FOR USE

*[Consult a physician for the following]*

1. The following persons should consult a physician, pharmacist, or registered distributor of this medication before using this medication.

A person with any serious symptoms, such as intense diarrhea, high fever, *etc.*

2. In the following cases, discontinue use immediately, and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

(1) Nervousness of children, night crying, peevishness, convulsion, loss of appetite, or gastrointestinal invalidism does not improve after having used this medication for a month.

(2) Common cold, fever caused by common cold, diarrhea, indigestion, or vomiting milk does not improve after having used this medication 5 or 6 times.

### INDICATIONS

Nervousness of children, night crying, peevishness, convulsion, common cold, fever caused by common cold, diarrhea, indigestion, vomiting milk, loss of appetite, gastrointestinal invalidism

### COMPOSITION (in 60 pills)

Components	Content
Agarwood	15.60 mg
Musk	0.78 mg
Ginseng	43.77 mg
Bear bile	0.90 mg

Additives: Rice Powder, Kanbaiko, *d*-Borneol, Honey, Paraben, Silver Leaf

#### DOSAGE AND ADMINISTRATION

Age	One dosage	Daily dosage
11~14 years	18~20 pills	3 times a day before or between each meal.
7~10 years	15~17 pills	
4~6 years	11~14 pills	
2~3 years	7~10 pills	
1 year	4~6 pills	
Under 1 year	1~3 pills	

#### *[Cautions]*

- (1) Please follow the recommended dosage and administration.
- (2) This medication should be given to children only under adult supervision.

#### CAUTIONS ON HANDLING AND STORAGE

- (1) Keep all medication out of the reach of children.
- (2) Do not transfer this medicine to a different container. (It may be misused or the quality may deteriorate.)
- (3) Store in a sealed container in a cool, dry place free from direct sunlight.