

*Please store this instruction leaflet in a safe place for later reference.*

*Be sure to read the instructions carefully before use.*

## **KIO FEVER (KODOMO GENETSU ZAYAKU)**

**キオフィーバ (こども解熱坐薬)**

<ANTIPYRETIC SUPPOSITORY>

“**KIO FEVER**” is antipyretic suppository for children. Approved for use by children aged 1 year up to 12 years.

### **CAUTIONS FOR USE**

(Though “**KIO FEVER**” is antipyretic for children, common cautions as antipyretic are described here.)

*[To be avoided!]*

(Failure to observe the following may result in the present symptoms worsening or adverse reactions/accidents occurring.)

1. This medication should not be used by the following persons:

- (1) Persons who experience an allergic reaction to this medication, and/or any component of this medication
- (2) Persons who suffer from asthma after using this medication or any other antipyretics or cold remedies

2. This medication should not be used in combination with the following medications:

Other antipyretics, cold remedies, and sedatives

3. Do not consume alcohol before or after taking this medication.

4. Do not use this medication for a prolonged period of time.

*[Consult a physician for the following]*

1. The following persons should consult a physician, dentist, pharmacist, or registered distributor of this medication before using this medication.

- (1) A person undergoing medical treatment from a physician or dentist
- (2) A pregnant woman or a woman suspected of being pregnant
- (3) An elderly person
- (4) A person with a medical history of being allergic to medicines and other substances
- (5) A person diagnosed as having the following:

Cardiac disease, renal disease, hepatic disease, gastric/duodenal ulcer

2. The following symptoms may be adverse reactions to this medication. If any of these

symptoms occur, discontinue use immediately and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

Related area	Symptoms
Skin	Rash, redness, itching
Digestive organs	Nausea, vomiting, loss of appetite
Central nervous system	Dizziness
Other	Significant decrease in temperature

In very rare case the following serious symptoms may occur. In this case, consult a physician immediately.

Symptom's name	Symptoms
Shock(anaphylaxis)	Occurrence of itching of skin, hives, hoarseness, sneezing, itching of throat, breathing difficulties, palpitations, or clouding of consciousness, etc. soon after use
Muco-cutaneo-ocular syndrome(Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalized exanthematous pustulosis	Continuation or sudden worsening of the following: high fever, redness of the eye, eye discharge, erosion of the lip, throat pain, rash or redness over an extensive area of skin, red circular skin bumps(pustules), general listlessness, loss of appetite, etc.
Hepatic function failure	Fever, itching, rash, jaundice(the skin and whites of the eyes turn yellow), brown urine, general listlessness, loss of appetite, etc.
Interstitial pneumonia	Sudden occurrence or continuation of the following: shortness of breath or choking sensation after going up stairs or following exertion, dry cough, fever, etc.
Asthma	Wheezing and labored breathing
Renal impairment	Decreased urinary output, fever, rash, decreased urinary output, general swelling, general listlessness, joint pain, diarrhea, etc.

- Discontinue use when the symptom does not improve after having used this medication once, and consult a physician, dentist, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

## INDICATIONS

Temporary relief of children's fever.

**COMPOSITION** (in 1 suppository)

Components	Content
Acetaminophen	100 mg

Additive: Hard Fat

**DOSAGE AND ADMINISTRATION**

Age	One dosage	Daily dosage
6 up to 12 years	1~2 suppositories	Insert into the rectum once a day.
3 up to 5 years	1 suppository	
1 up to 2 years	1/2~1 suppository	
Under 1 year	Do not use	

*[Cautions]*

- (1) This medication should be given to children only under adult supervision.
- (2) Please follow the recommended dosage and administration.
- (3) Do not use this medication to children under 1 year.
- (4) Use this medication only one day. Do not continue using for two straight days.
- (5) Do not take this medication orally.

**CAUTIONS ON HANDLING AND STORAGE**

- (1) Store in a sealed container in a cool (under 30 degree C), dry place free from direct sunlight (e. c. in the refrigerator).
- (2) Keep all medication out of the reach of children.
- (3) Do not transfer this medicine to a different container. (It may be misused or the quality may deteriorate.)